## **Suspected Concussion Report Form**



Player Name:				Playe	er DOB:				
Date & Time of Injury:									
Division:Level:	G	ame/Pr	actice Location:			Sex	: M	F	
Position during Injury (please circle):						Goa			
Injury Description: Player to player cont	act 🗆	]	Ball to player co	ntact□		Fall to grour	nd 🗆	Other 🗆	
Reported and Observable Symptoms (C	1								
Headache			entally foggy			ensitive to lig			
🗌 Nausea			owed down			ensitive to no	oise		
Dizziness		-	concentrating			rritability			_
□ Vomiting			remembering			adness			
Visual problems		rowsine	SS			lervous/anxio	ous		
Balance problems	□ S	leeping r	nore/less than us	ual		/lore emotion	al		
Numbness/Tingling	□ T	rouble fa	alling asleep		🗆 F	atigue			
Red Flag Symptoms (Check all that appl	ly): Ca	all 911 i	mmediately wit	h a su	dden ons	set of any of	these s	symptoms	
□Severe or increasing headache		🗆 Neck	pain or tenderness			Seizure or conv	ulsion		
Double vision		🗆 Loss	of consciousness		□ I	Repeated vomit	ting		
Weakness or tingling/burning in arms/legs		🗌 Dete	riorating conscious	state		ncreasingly res	stless, agi	tated or combative	
Are there any <u>other observable/report</u> If yes, what:	-	-		No					
Is there evidence of injury to anywhere If yes, where:			-	P □Y€	es 🗆 N	0			
Has this player had a concussion before If yes, how many:	e? [	] Yes		on't kno	ow 🗆	Prefer not t	to answ	ver	
Does this player have any pre-existing a lf yes, please list: Does this player take any medication?	□Yes	5 🗆 N	o 🗆 Don't kn	ow	□Prefer	not to answ		not to answer	
I [name of coach completing this form]: _								recommended	to
the player's parent or guardian that the j								-	2
					-		-		
family physician, pediatrician, sports-me Documentation from any other source w		• •		t or int	ternal me	edicine/reha	itati	on (physiatrists).	
Detamentation nom any other source w			<u>cptables</u>						
Signature	Date	e:		Ro	le:			_	
Phone Number:									

This form is to be completed by the head coach in the event of a <u>suspected concussion</u> in any club and/or team soccer activity. Once complete, give one copy of this report to parent/guardian and the other to <u>TSA office</u>. Parents must take this form to medical appointment with <u>medical doctor or nurse practitioner</u>. This report form is aligned with <u>best-practice</u> <u>guidelines</u> and a tool to be used to support the remove, refer and report sections of the <u>Canada Soccer Concussion Policy</u>.

	etimbling slow		Nausse of womiting       Fatigue of iow energy browines       Neck Pain ownicus       Feeling slowed down <ul> <li>Divisines</li> <li>Dowines</li> <li>Dowines</li></ul>	of the following signs are be safely and immediately be safely and immediately Deteriorating conscious state Vomiting Increasingly restless, agitated or combative agitated or combative or attempt to more the player than required for airway or of undess trained for airway and the equipment unless det to do so safely. Balance, gait difficulties,	the basic principles • 001 anget, response, (oth hing, circulation) • 001 for a spinal • 001 for a spinal rail critical. trail critical. trail sins of possible concussion should ible concussion include ible concussion include isorientation or • • • • • • • • • • • • • • • • • •
to respond appropriately stumbling, slow		apport sugges sugger or Not vot vot the CF the CF common	IMMEDIATELY REMOVED FROM PRACTICE OR PLAY AND SHOULD NOT RETURN TO ACTIVITY UNTIL ASSESSED MEDICALLY, EVEN	motor incoordination, stumbling, slow	onfusion, or an inability respond appropriately
motor incoordination.	motor incoordination,		ANY ATHLETE WITH A SUSPECTED CONCUSSION SHOULD BE	Balance, gait difficulties,	isorientation or
<ul> <li>Balance, gait difficulties, motor incoordination.</li> </ul>	Disorientation or     Balance, gait difficulties,     confusion, or an inability motor incoordination,     transmond anomorately etymological elow				sible concussion include
ggest possible concussion include:           • Disorientation or         • Balance, gait difficulties, confusion, or an inability	Balance, gait difficulties, motor incoordination, etumblinn elow		the Concussion in Sport vroup, it should not be altered in any way, repranoed or sold t commercial gain.		SNS
Balance, gait difficulties, motor incoordination.	Balance, gait difficulties, motor incoordination,	<ul> <li>* "Which half is it now?" * "Which half is it now?" * "Who scored last suggest a concussion:</li> <li>* "Who scored last in this game?"</li> <li>Athletes with suspected concussion should:</li> <li>Not be left alone initially (at least for the first 1-2 hours).</li> <li>Not drink alcohol.</li> <li>Not use recreational/ prescription drugs.</li> <li>Not use recreational/ prescription drugs.</li> <li>Not drive a motor vehicle until cleared to do so by a healthcare</li> </ul>	The CRT5 may be freely copied in its current form for distribution to individuals, teams, group and organisations. Any revision and any reproduction in a digital form requires approval the Concussion in Sport Group. It should not be altered in any way rebranded or sold f	proceed to the following steps:	of possible concussion should
proceed to the following steps: Balance, gait difficulties, motor incoordination.		<ul> <li>* "Which half is it now?"</li> <li>* "Who scored last sport) correctly may</li> <li>* "Who scored last in this game?"</li> <li>Athletes with suspected concussion should:</li> <li>Not be left alone initially (at least for the first 1-2 hours).</li> <li>Not use recreational/ prescription drugs.</li> <li>Not use recreational/ prescription drugs.</li> <li>Not be sent home by themselves. They need to be with a respo</li> </ul>	<ul> <li>Not drive a motor vehicle until cleared to do so by a healthcare professional.</li> </ul>	other equipment unces ed to do so safely.	
		<ul> <li>*Which half is it now?*</li> <li>*Who scored last sport) correctly may</li> <li>*Who scored last in this game?*</li> <li>Not be left alone initially (at least for the first 1-2 hours).</li> <li>Not use recreational/ prescription drugs.</li> </ul>	<ul> <li>Not be sent home by themselves. They need to be with a responsible adult.</li> </ul>	ot remove a helmet or other equipment unless	•
		<ul> <li>* "Which half is it now?"</li> <li>* "Who scored last suggest a concussion:         <ul> <li>* "Who scored last in this game?"</li> <li>Athletes with suspected concussion should:</li> <li>Not be left alone initially (at least for the first 1-2 hours).</li> <li>Not the recreational/ mesorintion druce</li> </ul> </li> </ul>		port) unless trained to so do.	
		<ul> <li>*Which half is it now?*</li> <li>*Which half is it now?*</li> <li>suggest a concussion:         <ul> <li>*Who scored last in this game?*</li> </ul> </li> <li>Athletes with suspected concussion should:         <ul> <li>Not be left alone initially (at least for the first 1-2 hours).</li> <li>Not drink alcohol.</li> </ul> </li> </ul>	<ul> <li>Not use recreational/ prescription drugs.</li> </ul>	or attempt to move the project or than required for airway	
		<ul> <li>*Which half is it now?*</li> <li>*Which half is it now?*</li> <li>suggest a concussion:         <ul> <li>*Who scored last in this game?*</li> </ul> </li> <li>Athletes with suspected concussion should:         <ul> <li>Not be left alone initially (at least for the first 1-2 hours).</li> </ul> </li> </ul>	Not drink alcohol.	of attempt to move the player	•
		<ul> <li>appropriately for each sport or reach suggest a concussion:</li> <li>"Who scored last in this game?"</li> <li>Athletes with suspected concussion should:</li> </ul>	<ul> <li>Not be left alone initially (at least for the first 1-2 hours).</li> </ul>		
		<ul> <li>appropriately for each</li> <li>"Which half is it now?"</li> <li>suggest a concussion:</li> <li>"Who scored last in this game?"</li> </ul>	Athletes with suspected concussion should:	Increasingly restless, agitated or combative	
			<ul> <li>"Which half is it now?"</li> <li>"Who scored last in this game?"</li> </ul>	e professional is available, Deteriorating Conscious state Vomitinn	Loss of consciousness •
			(IN ATHLETES OLDER THAN 12 YEARS)		ry including whether ANY ried then the player should y. If no licensed healthcare dical assessment: Severe or increasing headache Seizure or convulsion Loss of consciousness
		ANCE	STEP 4: MEMORY ASSESSMENT		AN AMBULANCE ry including whether ANY rted then the player should dy. If no licensed healthcare dical assessment: Sever or increasing • headache Seizure or convulsion • Loss of consciousness •
			low energy • Neck Pain • "Don't feel right"	es. The Concussion Recognition Tool t designed to diagnose concussion.	and potentially fatal brain injurie of suspected concussion. It is no AN AMBULANCE ry including whether ANY rised then the player should dical assessment: Gevere or increasing headache Seizure or convulsion Loss of consciousness
		<ul> <li>Drowsiness low energy . Neck Pain</li> <li>Dizziness . "Don't feel right" .</li> <li>Dizziness . "Don't feel right" .</li> <li>In ziness</li></ul>	Fatigue or anxious	<ol> <li>The Concussion Recognition Tool t designed to diagnose concussion.</li> </ol>	of suspected concussion. It is no of suspected concussion. It is no AN AMBULANCE Try including whether ANY vited then the player should vy. If no licensed healthcard dical assessment: Severe or increasing headache Seizure or convulsion Loss of consciousness
		<ul> <li>Balance problems · Sensitivity · Sadness to noise to noise · Nervous or vomiting · Fatigue or anxious low energy · Neck Pain · Drowsiness · "Don't feel right" · Neck Pain · Dizziness · "Don't feel right" · Neck Pain · Nec</li></ul>	<ul> <li>Sensitivity</li> <li>Sadness</li> <li>to noise</li> </ul>	A.The Concussion Recognition Tool t designed to diagnose concussion.	Supported by and potentially fatal brain injurit of suspected concussion. It is no AN AMBULANCE AN AMBULANCE AN AMBULANCE An end the player and dical assessment: Severe or increasing headache Seizure or convulsion Loss of consciousness
		<ul> <li>'Pressure in head"</li> <li>Balance problems</li> <li>Balance problems</li> <li>Sensitivity</li> <li>Sensitivity</li> <li>Sadness</li> <li>Nervous or anxious</li> <li>Drowsiness</li> <li>Drowsiness</li> <li>'Don't feel right"</li> <li>Dizziness</li> <li>Nature or anxious</li> <li>Neck Pain</li> <li>Natious</li> <li>Neck Pain</li> <li>Natious</li> <li>Neck Pain</li> <li>Natious</li> </ul>	<ul> <li>Sensitivity to light</li> <li>More Irritable</li> <li>Sensitivity</li> <li>Sadness</li> <li>to noise</li> </ul>	a: The Concussion Recognition Tool	Supported by Supported by and potentially fatal brain injurit of suspected concustion. It is no any including whether ANY ried then the player should y. If no licensed healthcard dical assessment: Severe or increasing headache headache Loss of consciousness
		<ul> <li>Headache</li> <li>Headache</li> <li>Blurred vision</li> <li>More emotional</li> <li>"Pressure in head"</li> <li>Sensitivity to light</li> <li>More Irritable</li> <li>Balance problems</li> <li>Sensitivity</li> <li>Sensitivity</li> <li>Sensitivity</li> <li>Sensitivity</li> <li>Nervous or</li> <li>Nervo</li></ul>	Blurred vision     More emotional     In head     Sensitivity to light     More Irritable     roblems     Sensitivity     sadness     to noise	escents and adults	sion in children, adol Supported by and potentially fatal brain injurit of suspected concussion. It is no a NAMBULANCE AN AMBULANCE AN AMBULANCE AN ambulance assessment: Severe or increasing headache headache toss of consciousness